Solar activity ranged from very low to low. Low levels occurred on 23 February while very low conditions were observed on 21 – 22 and 24 - 27 February. The largest region on the disk, Region 735 (S07, L=121, class/area, Ekc/470 on 16 February), remained fairly quiet, producing only minor activity before rotating around the west limb on 23 February. Midday on 27 February, a section of a large filament erupted off the southwest limb. LASCO imagery indicated an associated CME was observed off the southwest limb; however, it did not appear to be Earth directed. All other regions on the disk were magnetically simple and stable.

Solar wind data were available from the NASA Advanced Composition Explorer (ACE) spacecraft during most of the summary period. The period began with solar wind speed near 425 km/s and the Bz component of the IMF weak, not varying much beyond +/- 4 nT. Thereafter, and through early on 23 February, the IMF Bz remained neutral while the solar wind speed hit a low of near 340 km/s. On 23 February, a solar sector boundary crossing was observed in the ACE satellite data. By late on the 23rd, the IMF Bz began to oscillate through +/- 11 nT while temperature, density, and solar wind speed increased, all indicative of a co-rotating interactive region (CIR) and a high speed coronal hole wind stream. Wind speed gradually increased, and ended the summary period near 615 km/s while the IMF Bz remained variable, not fluctuating much beyond +/-10 nT. For the balance of the summary period, the geomagnetic field remained under the influence of the high speed stream.

No greater than 10 MeV proton events were observed this period.

The greater than 2 MeV electron flux at geosynchronous orbit was at high levels from 21-24 February.

During the summary period, the geomagnetic field ranged from quiet to unsettled levels with active to minor storming periods observed at the higher latitudes. The period began on 21 February with quiet to unsettled conditions with periods of active conditions observed at the higher latitudes. From 22 February through midday on the 24th, conditions were quiet at all latitudes. By midday on the 24th, activity increased to quiet to unsettled with periods of active to minor storming observed at high latitudes due to the effects of a geoeffective coronal hole high speed wind stream. These conditions persisted for the remainder of the summary period.

Space Weather Outlook 02 March - 28 March 2005

Solar activity is expected be at very low to low conditions the entire forecast period.

A greater than 10 MeV proton event is not expected.

The greater than 2 MeV electron flux at geosynchronous orbit is expected to be at high levels on 07 – 12 March and again 17 – 19 March.

The geomagnetic field is expected to range from quiet to minor storm levels. Coronal hole high speed wind streams are expected to produce unsettled to active levels with occasional minor storm periods on 06 - 09 March and again on 16 - 17 March. Otherwise, expect quiet to unsettled conditions.



Daily Solar Data

				Duny So	uu D	uu						
	Radio	Sun	Sunspot	X-ray	_			Flares				
	Flux	spot	Area	Background	X	-ray F	lux		Op	otical		
Date	10.7 cm	No.	(10 ⁻⁶ hemi.)		С	M	X	S	1	2	3	4
21 February	95	33	560	A9.9	0	0	0	0	0	0	0	0
22 February	92	23	510	B1.1	0	0	0	0	0	0	0	0
23 February	85	54	200	B1.1	2	0	0	0	0	0	0	0
24 February	80	17	40	A6.4	0	0	0	0	0	0	0	0
25 February	78	15	30	A2.1	0	0	0	0	0	0	0	0
26 February	77	27	70	A1.4	0	0	0	0	0	0	0	0
27 February	76	12	20	A1.7	0	0	0	0	0	0	0	0

Daily Particle Data

		oton Fluence ons/cm ² -day-s	r)	Electron Fluence (electrons/cm²-day-sr)
Date	>1MeV	>10MeV	>100MeV	>.6MeV >2MeV >4MeV
21 February	2.7E+5	1.3E+4	2.8E+3	3.9E+7
22 February	3.2E + 5	1.3E+4	2.9E + 3	6.9E+7
23 February	3.5E+5	1.4E+4	3.1E+3	9.9E+7
24 February	5.0E+5	1.4E+4	3.2E + 3	7.3E+7
25 February	3.9E + 5	1.4E+4	3.2E + 3	7.5E+6
26 February	3.0E+5	1.4E+4	3.5E+3	3.8E+6
27 February	3.7E+5	1.4E+4	3.4E+3	8.3E+6

Daily Geomagnetic Data

		Middle Latitude High Latitude				Estimated
		Fredericksburg		College		Planetary
Date	Α	A K-indices		K-indices	A	K-indices
21 February	4	3-2-2-0-1-0-0-1	18	2-3-3-4-4-2-4-3	8	3-3-1-2-1-2-1-2
22 February	3	1-1-0-0-1-1-2-2	2	0-0-0-0-0-2-2-1	4	1-1-0-0-1-2-2-1
23 February	2	1-0-1-1-1-0-1-1	1	0-0-1-0-0-0-1	4	1-1-2-1-1-2-1-1
24 February	4	0-1-1-1-3-1-1-1	8	0-0-0-1-5-1-2-1	5	0-0-1-1-3-1-2-2
25 February	9	1-1-2-2-3-3-3-2	12	0-0-2-3-3-4-4-1	9	1-1-2-2-3-3-2
26 February	9	2-2-2-3-3-2-1	18	1-1-2-4-5-5-2-1	9	2-2-2-3-3-3-2-2
27 February	6	0-2-2-1-3-2-1-1	9	0-1-2-2-5-1-1-1	8	0-2-3-1-3-2-2-1




Alerts and Warnings Issued

Date & Time of Issue	Type of Alert or Warning	Date & Time of Event UTC
21 Feb 0006	1 – 245 MHz Radio Noise Storm	20 Feb
21 Feb 1436	ALERT: Electron 2MeV Integral Flux > 1000pfu	21 Feb 1415
22 Feb 0006	1 – 245 MHz Radio Noise Storm	21 Feb
22 Feb 1513	ALERT: Electron 2MeV Integral Flux > 1000pfu	22 Feb 1450
23 Feb 1157	ALERT: Electron 2MeV Integral Flux > 1000pfu	22 Feb 1140
24 Feb 0018	4 – 245 MHz Radio Bursts	23 Feb
24 Feb 1114	ALERT: Electron 2MeV Integral Flux > 1000pfu	24 Feb 1055
25 Feb 1706	WARNING: Geomagnetic K= 4	25 Feb 1707 - 2359
25 Feb 1739	ALERT: Geomagnetic K=4	25 Feb 1738
25 Feb 2353	EXTENDED WARNING: Geomagnetic K= 4	25 Feb 1707 -26 Feb 1600
26 Feb 1555	EXTENDED WARNING: Geomagnetic K= 4	25 Feb 1707 -27 Feb 1600



Twenty-seven Day Outlook



	Radio Flux	Planetary	Largest		Radio Flux	Planetary	Largest
Date	10.7 cm	A Index	Kp Index	Date	10.7 cm	A Index	Kp Index
02 Mar	75	8	3	16 Mar	100	15	4
03	80	8	3	17	100	25	5
04	80	5	2	18	95	12	3
05	85	5	2	19	90	10	3
06	85	15	4	20	90	5	2
07	90	20	5	21	90	5	2
08	90	20	5	22	85	5	2
09	95	15	4	23	80	5	2
10	100	10	3	24	75	10	3
11	100	10	3	25	75	10	3
12	100	8	3	26	75	10	3
13	100	10	3	27	75	10	3
14	100	8	3	28	75	12	3
15	100	8	3				



Energetic Events

				Birei Sei	ic Brents					
	Time		X-ray	Opt	ical Information	1	Peak	Sweep Freq		
Date		1/2	Integ	Imp/	Location	Rgn	Radio Flux	Intensity		
	Begin Ma	Max	Class Flux	Brtns	Lat CMD	#	245 2695	II IV		
	No Events	Observ	red .							

Flare List

				Flare List			
						ptical	
Data	Davis	Time	E. J	X-ray	Imp /	Location	Rgn
Date	Begin	Max	End	Class.	Brtns	Lat CMD	
21 February	0852	0856	0900	B1.8			
	1044	1048	1058	B1.7			
	1234	1238	1240	B2.2			735
	1331	1334	1336	B1.4			736
	1416	1422	1425	B6.4			736
	1957	2000	2006	B2.8			736
	2014	2019	2028	B6.0			736
	2212	2216	2220	B1.9			
	2241	2245	2248	B2.0			735
22 February	1409	1414	1424	B2.6			734
	1624	1628	1637	B2.1			736
	1806	1859	1936	B8.9			
	2115	2119	2122	B4.1			
23 February	0540	0544	0546	B2.6			
	0559	0603	0605	B2.2			735
	0621	0628	0633	C1.3			735
	0731	0744	0751	B1.8			735
	0815	0823	0827	B2.4			736
	0844	0848	0850	B8.1			736
	0931	0934	0936	C2.6			736
	1144	1148	1150	B2.6			736
	1359	1404	1406	B6.0			736
24 February	0007	0022	0038	B2.4			737
25 February	No Fla	resObserv	ved				
26 February	No Fla	resObserv	ved				
27 February	No Fla	resObserv	ved				



Region Summary

	Locatio	n			Characte		<u> </u>			F	lares					
	Locatio	Helio	Area	Extent	Spot	Spot	Mag		X-ra		iarc		Ontic	al	_	
Date (° Lat °	CMD)	Lon	(10 ⁻⁶ hemi	(helio)	Class	Count	Class	C	M	X	S	1	2	3	4	
	Re	gion 73	24													
09 Feb S04I		136	0060	02	Hax	001	A									
10 Feb S04I		137	0050	02	Hax	001	A	1								
11 Feb S04I		136	0060	02	Hsx	001	A	1								
12 Feb S05H		136	0060	02	Hsx	001	A									
13 Feb S04I	E1 7	137	0080	02	Hsx	001	A	1			1					
14 Feb S05F	E 04	137	0070	05	Cao	007	В	1			1					
15 Feb S04V	W10	138	0070	04	Cso	004	В	1								
16 Feb S04V	W23	138	0080	04	Cso	005	В	1				1				
17 Feb S05V	W34	135	0070	04	Cso	004	В									
18 Feb S05V	W50	138	0040	02	Dso	003	В									
19 Feb S05V	W63	138	0040	01	Hsx	001	Α									
20 Feb S05V	W76	138	0040	01	Hsx	001	Α									
21 Feb S07V	W89	138														
								6	0	0	2	1	0	0	0	
Crossed Wes																
Absolute hel	iogra	ohic lon	gitude:137													
	Re	gion 73	25													
10 Feb S08I		120	0080	04	Hsx	001	A									
11 Feb S08I	E60	120	0410	06	Dho	003	В									
12 Feb S08F	E 47	120	0380	04	Hhx	001	A				1					
13 Feb S09I	E 34	120	0370	07	Dho	008	В									
14 Feb S08I	E 20	121	0440	07	Dko	026	В									
15 Feb S08I	E06	122	0420	11	Cko	015	Bg				3					
16 Feb S07V	W06	121	0470	11	Ekc	017	Bgd	1			1					
17 Feb S10V	W18	119	0420	08	Dkc	011	Bgd									
18 Feb S09V		120	0460	06	Dko	008	В	1			1					
19 Feb S09V		120	0410	06	Dki	018	Bg									
20 Feb S09V		120	0440	06	Dko	012	Bg									
21 Feb S09V		120	0440	05	Dko	006	Bg									
22 Feb S09V		120	0300	03	Cko	002	В									
23 Feb S09V	W97	119	0060	02	Hsx	001	A	1								
								3	0	0	6	0	0	0	0	
Crossed Wes	at I im	h														

Crossed West Limb. Absolute heliographic longitude:122



Region Summary – continued.

			Re	egion Si			<u>tinued.</u>									
	Locatio		Sunspot Characteristics						Flares							
Data	(01 at 0 CMD)	Helio	Area (10 ⁻⁶ hemi)	Extent	Spot	Spot	Mag	$\overline{\mathbf{C}}$	X-ra	_		1	Optic 2			
_Date	(°Lat°CMD)	Lon	(10 nemi)	(nello)	Class	Count	Class		M	X	S	_1_		3	4	
	Re	gion 730	5													
20 Feb	N13W50	112	0030	05	Dao	006	В									
21 Feb	N13W63	112	0120	07	Dao	007	В									
22 Feb	N13W77	112	0210	07	Dko	001	В									
23 Feb	N13W92	114	0090	02	Cso	003	В	1								
								1	0	0	0	0	0	0	0	
Crosse	ed West Lim	ıb.														
Absol	ute heliograj	phic long	gitude:112													
	$R_{\mathcal{O}}$	gion 737	7													
23 Fel	S07W31	053	0040	04	Dsi	008	В									
	S09W44	053	0040	05	Dso	007	В									
	S11W59	055	0030	06	Bxo	005	В									
	S11W72	055	0040	04	Bxo	003	В									
	S11W72	055	0040	04	DAU	003	Ъ									
2/100) 511 W 05	033						0	0	0	0	0	0	Λ	0	
Still o	n Disk.							U	U	U	U	U	U	U	U	
	ute heliograj	nhic lone	ritude: 053													
AUSUI		_														
		gion 738														
	S10E10	012	0010	02	Bxo	002	В									
24 Fel	S10W03	012														
								0	0	0	0	0	0	0	0	
	n Disk.															
Absol	ute heliogra _l	phic long	gitude: 012													
	Re	gion 739	9													
26 Fel	S04E52	291	0030	05	Bxo	004	В									
27 Fel	S04E38	292	0020	01	Axx	002	A									
•								0	0	0	0	0	0	0	0	
Still o	n Disk.							-	-	-	-	-	-	-	-	
	ute heliogra	phic long	gitude: 292													
	- 31		–													




Recent Solar Indices (preliminary)
of the observed monthly mean values

_				of the o	<u>observed n</u>	nonthly i	<u>mean values</u>			
			Sunsp	ot Number	Radio	Flux	Geomagnetic			
		Observed	values	<u>Ratio</u>	Smooth	values	*Penticton	Smooth	Planetary	Smooth
	Month	SWO	RI	RI/SWO	SWO	RI	10.7 cm	Value	Ap	Value
					,	2003				
	February	87.0	46.0	0.53	136.4	78.5	124.5	144.7	17	18.9
	March	119.7	61.1	0.51	128.1	74.2	132.2	139.5	21	19.4
	TVIGICII	117.7	01.1	0.51	120.1	7 1.2	132.2	137.3	21	17.1
	April	119.7	60.0	0.50	121.5	70.3	126.3	135.0	20	20.1
	May	89.6	55.2	0.62	118.3	67.8	129.3	133.1	26	21.0
	June	118.4	77.4	0.65	113.6	65.2	129.4	130.2	24	21.5
	July	132.8	85.0	0.64	106.9	62.0	127.8	127.2	19	22.0
	August	114.3	72.7	0.64	102.8	60.3	122.1	125.2	23	22.2
	September	82.6	48.8	0.59	100.7	59.8	112.3	123.7	18	21.8
	October	118.9	65.5	0.55	96.6	58.4	153.1	121.8	35	21.1
	November	118.9	67.3	0.57	93.6	57.0	153.1	120.1	28	20.0
	December	75.4	46.5	0.62	91.4	55.0	115.1	118.0	16	18.6
					,	2004				
	January	62.3	37.7	0.61	87.9	52.0	114.1	116.3	22	18.1
	February	75.6	45.8	0.61	84.2	49.4	107.0	115.5	13	17.7
	March	81.0	49.1	0.61	80.9	47.2	112.2	114.6	14	16.9
	April	59.3	39.3	0.66	77.9	45.6	101.2	112.3	11	15.5
	May	77.3	41.5	0.54	74.1	43.9	99.8	109.2	8	14.3
	June	78.9	43.2	0.55	70.4	41.7	97.4	107.2	8	14.0
	July	87.8	51.0	0.58	68.3	40.2	118.5	105.9	23	13.8
	August	69.5	40.9	0.59			110.1		11	
	September	50.0	27.7	0.55			103.1		10	
	October	77.9	48.4	0.62			105.7		9	
	November		43.7	0.62			113.2		26	
	December	34.7	17.9	0.52			94.6		11	
	December	57.7	11.7	0.52		•••	74.0		11	
	T	52 0	21.2	0.60	2	2005	100 4		22	
	January	52.0	31.3	0.60			102.4		22	

NOTE: All smoothed values after September 2002 and monthly values after March 2003 are preliminary estimates. The lowest smoothed sunspot index number for Cycle 22, RI = 8.0, occurred in May 1996. The highest smoothed sunspot number for Cycle 23, RI= 120.8, occurred April 2000. *After June 1991, the 10.7 cm radio flux data source is Penticton, B.C. Canada. Prior to that, it was Ottawa.





Weekly Geosynchronous Satellite Environment Summary Week Beginning 21 February 2005

Protons plot contains the five-minute averaged integral proton flux (protons/cm²-sec -sr) as measured by GOES-11 (W110) for each of three energy thresholds: greater than 10, 50, and 100 MeV.


Electrons plot contains the five-minute averaged integral electron flux (electrons/cm² –sec –sr) with energies greater than 2 MeV at GOES-12 (W75).

Hp plot contains the five minute averaged magnetic field H - component in nanoteslas (nT) as measured by GOES-12. The H component is parallel to the spin axis of the satellite, which is nearly parallel to the Earth's rotation axis.

Kp plot contains the estimated planetary 3-hour K-index (derived by the Air Force Weather Agency) in real time from magnetometers at Meanook, Canada; Sitka, AK; Glenlea, Canada; St. Johns, Canada; Ottawa, Canada; Newport, WA; Fredericksburg, VA; Boulder, CO; Fresno, CA and Hartland, UK. These data are made available through cooperation from the Geological Survey of Canada (GSC), British Geological Survey (BGS) and the US Geological Survey. These may differ from the final Kp values derived from a more extensive network of magnetometers.

The data included here are those now available in real time at the SWO and are incomplete in that they do not include the full set of parameters and energy ranges known to cause satellite operating anomalies. The proton and electron fluxes and Kp are "global" parameters that are applicable to a first order approximation over large areas. H parallel is subject to more localized phenomena and the measurements generally are applicable to within a few degrees of longitude of the measuring satellite.





Weekly GOES Satellite X-ray and Proton Plots

X-ray plot contains five-minute averaged x-ray flux (watts/m²⁾ as measured by GOES 12 (W76) and GOES 10 (W135) in two wavelength bands, .05 - . 4 and .1 - .8 nm. The letters A, B, C, M and X refer to x-ray event levels for the .1 - .8 nm band.

Proton plot contains the five-minute averaged integral proton flux (protons/cm² –sec-sr) as measured by GOES-11 (W110) for each of the energy thresholds: >1, >10, >30 and >100 MeV. P10 event threshold is 10 pfu (protons/cm²-sec-sr) at greater than 10 MeV.

